

# Ionization Rates for 1963-2005 from Solar Proton Events

Charles H. Jackman

E-mail: Charles.H.Jackman@nasa.gov

Phone: 301-614-6053

Code 613.3

Laboratory for Atmospheres

NASA Goddard Space Flight Center

Greenbelt, MD 20771

October 2006

**Abstract:** Large solar eruptive occurrences can lead to significant fluxes of protons at the Earth, which are called solar proton events (SPEs). Proton flux measurements during SPEs from satellites over the period 1963-2005 have been used to compute daily average ion pair production rates using an energy deposition calculation described below. These daily average ion pair production rates as functions of pressure between 888 hPa (~1 km) and  $8 \times 10^{-5}$  hPa (~115 km) are provided at the SOLARIS website (<http://strat-www.met.fu-berlin.de/~matthes/sparc/inputdata.html>) and can be assumed to affect the atmosphere approximately uniformly over both polar cap regions (60-90°N and 60-90°S geomagnetic latitude). Methodologies for deriving HO<sub>x</sub> (H, OH, HO<sub>2</sub>) and NO<sub>x</sub> (N, NO, NO<sub>2</sub>) production rates from these ion pair production rates are also given below.

## 1. Proton Flux and Ion Pair Production

Solar proton fluxes have been measured by a number of satellites in interplanetary space or in orbit around the Earth. The National Aeronautics and Space Administration (NASA) Interplanetary Monitoring Platform (IMP) series of satellites provided measurements of proton fluxes from 1963-1993. IMPs 1-7 were used for the fluxes from 1963-1973 [Jackman et al. 1990] and IMP 8 was used for the fluxes from 1974-1993 [Vitt and Jackman, 1996]. The National Oceanic and Atmospheric Administration (NOAA) Geostationary Operational Environmental Satellites (GOES) were used for proton fluxes from 1994-2005 [Jackman et al. 2005a].

Proton flux data were taken from T. Armstrong and colleagues (University of Kansas, private communication, 1986) for the period 1963-1973 (see Armstrong et al. 1983 for a discussion of the IMP 1-7 satellite measurements). These data were fit with a power law form that was assumed to be valid over the range 5-100 MeV [Jackman et al. 1990] and then degraded in energy using the scheme first discussed in Jackman et al. [1980]. This energy degradation scheme assumes that the energy lost in each of the atmospheric slabs can be quantified using well-known range-energy relationships from

Sternheimer [1959]. The scheme divides the protons into 60 monoenergetic energy intervals, all assumed to be isotropic, as well as 35 pitch angles. The scheme includes the deposition of energy by all the protons and associated secondary electrons. The energy required to create 1 ion pair was assumed to be 35 eV [Porter et al. 1976]. This energy degradation scheme has been compared with two other independent methodologies [Lummerzheim, 1992] and is found to be in good agreement with these other schemes.

IMP 8 was used for the proton flux data for the years 1974-1993. Vitt and Jackman [1996] take advantage of the measurements of alpha particles by IMP 8 as well and use proton fluxes from 0.38-289 MeV and alpha fluxes from 0.82-37.4 MeV in energy deposition computations. The energy deposition methodology is similar to that discussed in Jackman et al. [1980]. Alpha particles were found to add about 10% to the total ion pair production during SPEs.

Four GOES satellites are used for the proton fluxes in years 1994-2005: 1) GOES-7 for the period January 1, 1994 through February 28, 1995; 2) GOES-8 for the period March 1, 1995 through April 8, 2003, and May 10, 2003 to June 18, 2003; 3) GOES-11 for the period June 19, 2003 to December 31, 2005; and 4) GOES-10 to fill in the gap of missing proton flux data from April 9 through May 9, 2003. The GOES satellite proton fluxes are fit in three energy intervals 1-10 MeV, 10-50 MeV, and 50-300 MeV with exponential spectral forms. The energy deposition methodology again is that discussed in Jackman et al. [1980].

<b>Table 1.</b> List of years, publication reference, and source of proton fluxes used in creating the ion pair production rates.		
Years	Reference	Source of Protons
1963-1973	Jackman et al. [1990]	IMP 1-7
1974-1993	Vitt and Jackman [1996]	IMP 8
1994-2005	Jackman et al. [2005a]	GOES 7, 8, 10, 11

**The daily average ion pair production rates are provided as functions of pressure between 888 hPa (~1 km) and  $8 \times 10^{-5}$  hPa (~115 km) at the SOLARIS website (<http://strat-www.met.fu-berlin.de/~matthes/sparc/inputdata.html>) and can be assumed to affect the atmosphere approximately uniformly over both polar cap regions (60-90°N and 60-90°S geomagnetic latitude). The datasets are divided into 43 individual yearly ASCII files labeled: *IonPair\_Year\_1963.dat...* *IonPair\_Year\_2005.dat* A fortran reader is provided labeled: *IP\_read.f***

## 2. Odd Hydrogen (HO<sub>x</sub>) Production

Along with the ion pairs, the protons and their associated secondary electrons also produce odd hydrogen (HO<sub>x</sub>) and odd nitrogen (NO<sub>y</sub>). The production of HO<sub>x</sub> relies on complicated ion chemistry that takes place after the initial formation of ion pairs [Swider and Keneshea, 1973; Frederick, 1976; Solomon et al. 1981]. Solomon et al. [1981] computed HO<sub>x</sub> production rates as a function of altitude and ion pair production. Some of these computations are given in Table 2 for background ion pair production rates of 10<sup>2</sup>, 10<sup>3</sup>, and 10<sup>4</sup> cm<sup>-3</sup>s<sup>-1</sup>. Each ion pair typically results in the production of around two HO<sub>x</sub> constituents in the upper stratosphere and lower mesosphere. In the middle and upper mesosphere, an ion pair is computed to produce less than two HO<sub>x</sub> constituents per ion pair.

**Table 2.** HO<sub>x</sub> constituents produced per ion pair as a function of altitude for baseline ionization rates (BIR) of 10<sup>2</sup>, 10<sup>3</sup>, and 10<sup>4</sup> cm<sup>-3</sup>s<sup>-1</sup>. These HO<sub>x</sub> production rates were taken from Solomon et al. [1981].

Altitude (km)	HO <sub>x</sub> production per ion pair		
	BIR – 10 <sup>2</sup> cm <sup>-3</sup> s <sup>-1</sup>	BIR – 10 <sup>3</sup> cm <sup>-3</sup> s <sup>-1</sup>	BIR – 10 <sup>4</sup> cm <sup>-3</sup> s <sup>-1</sup>
40	2.00	2.00	1.99
45	2.00	1.99	1.99
50	1.99	1.99	1.98
55	1.99	1.98	1.97
60	1.98	1.97	1.94
65	1.98	1.94	1.87
70	1.94	1.87	1.77
75	1.84	1.73	1.60
80	1.40	1.20	0.95
85	0.15	0.10	0.00
90	0.00	0.00	0.00

**The SPE-produced HO<sub>x</sub> constituents can be included in model simulations in the following way: Use an ASCII dataset (*HOx\_Prod\_Solomon.dat*) with a fortran reader (*HOx\_Prod\_Solomon.f*), which uses the Solomon et al. [1981] methodology in table form. These are located at the SOLARIS website: <http://strat-www.met.fu-berlin.de/~matthes/sparc/inputdata.html>**

### 3. Odd Nitrogen (NO<sub>y</sub>) Production

Odd nitrogen is produced when the energetic charged particles (protons and associated secondary electrons) collide with and dissociate N<sub>2</sub>. Following Porter et al. [1976] it is assumed that ~1.25 N atoms are produced per ion pair. The Porter et al. [1976] study also further divided the proton impact of N atom production between ground state (~45% or ~0.55 per ion pair) and excited state (~55% or ~0.7 per ion pair) nitrogen atoms. Ground state [N(<sup>4</sup>S)] nitrogen atoms can create other NO<sub>y</sub> constituents, such as NO, through



or can lead to NO<sub>y</sub> destruction through



Generally, excited states of atomic nitrogen, such as N(<sup>2</sup>D), result in the production of NO through



[e.g., Rusch et al., 1981; Rees, 1989] and do not cause significant destruction of NO<sub>y</sub>. Rusch et al. [1981] showed that there are huge differences in the final results of model computations of NO<sub>y</sub> enhancements from SPEs that depend strongly on the branching ratios of the N atoms produced. If a model does not include any of the excited states of atomic nitrogen [e.g., N(<sup>2</sup>D), N(<sup>2</sup>P), and N<sup>+</sup>] in their computations, the NO<sub>y</sub> production from SPEs can still be included.

**Here is a fairly accurate way to best represent the production of NO<sub>y</sub> constituents by the protons and their associated secondary electrons [Jackman et al. 2005b]: Assume that 45% of the N atoms produced per ion pair result in the production of N(<sup>4</sup>S) {~0.55 per ion pair} and that 55% of the N atoms produced per ion pair result in the production of NO {~0.7 per ion pair}.**

**Acknowledgments.** I thank Francis Vitt (NCAR) and Miriam Sinnhuber (for help in creating this ionization rate data set. I also thank the IMP and NOAA GOES teams for providing the solar proton flux data.

## References

- Armstrong, T. P., C. Brundardt, and J. E. Meyer, Satellite observations of interplanetary and polar cap solar particle fluxes from 1963 to the present, in *Weather and Climate Response to Solar Variations*, edited by B. M. McCormac, pp. 71-79, Colorado, Associated University Press, Boulder, 1983.
- Frederick, J. E., Solar corpuscular emission and neutral chemistry in the Earth's middle atmosphere. *J. Geophys. Res.*, *81*, 3179-3186, 1976.
- Jackman, C. H., J. E. Frederick, and R. S. Stolarski, Production of odd nitrogen in the stratosphere and mesosphere: An intercomparison of source strengths, *J. Geophys. Res.*, *85*, 7495-7505, 1980.
- Jackman, C. H., A. R. Douglass, R. B. Rood, R. D. McPeters, and P. E. Meade, Effect of solar proton events on the middle atmosphere during the past two solar cycles as computed using a two-dimensional model, *J. Geophys. Res.*, *95*, 7417-7428, 1990.
- Jackman, C. H., M. T. DeLand, G. J. Labow, E. L. Fleming, D. K. Weisenstein, M. K. W. Ko, M. Sinnhuber, J. Anderson, and J. M. Russell, The influence of the several very large solar events in years 2000-2003 on the neutral middle atmosphere, *Adv. Space Res.*, *35*, 445-450, 2005a.
- Jackman, C. H., M. T. DeLand, G. J. Labow, E. L. Fleming, D. K. Weisenstein, M. K. W. Ko, M. Sinnhuber, and James M. Russell, Neutral atmospheric influences of the solar proton events in October-November 2003, *J. Geophys. Res.*, *110*, A09S27, doi:10.1029/2004JA010888, 2005b.
- Lummerzheim, D., Comparison of energy dissipation functions for high energy auroral electron and ion precipitation, Geophysical Institute Report UAG-R-318, Geophysical Institute, University of Alaska Fairbanks, 1992.
- Porter, H. S., C. H. Jackman, and A. E. S. Green, Efficiencies for production of atomic nitrogen and oxygen by relativistic proton impact in air, *J. Chem. Phys.*, *65*, 154-167, 1976.
- Rees, M. H., Physics and chemistry of the upper atmosphere, pp. 278-281, Cambridge University Press, Cambridge, 1989.
- Rusch, D. W., J.-C. Gerard, S. Solomon, P. J. Crutzen, and G. C. Reid, The effect of particle precipitation events on the neutral and ion chemistry of the middle atmosphere, 1. Odd nitrogen, *Planet. Space Sci.*, *29*, 767-774, 1981.
- Solomon, S., D. W. Rusch, J.-C. Gerard, G. C. Reid, and P. J. Crutzen, The effect of particle precipitation events on the neutral and ion chemistry of the middle atmosphere, 2. Odd hydrogen, *Planet. Space Sci.*, *29*, 885-892, 1981.
- Sternheimer, R. M., Range-energy relations for protons in Be, C, Al, Cu, Pb, and air, *Phys. Rev.*, *115*, 137-142, 1959.
- Swider, W., and T. J. Keneshea, Decrease of ozone and atomic oxygen in the lower mesosphere during a PCA event, *Planet. Space Sci.*, *21*, 1969-1973, 1973.
- Vitt, F. M., and C. H. Jackman, A comparison of sources of odd nitrogen production from 1974 through 1993 in the Earth's middle atmosphere as calculated using a two-dimensional model, *J. Geophys. Res.*, *101*, 6729-6739, 1996.